

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2005/007052

International filing date: 08 March 2005 (08.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04380057.2
Filing date: 12 March 2004 (12.03.2004)

Date of receipt at the International Bureau: 12 December 2006 (12.12.2006)

Remark: Priority document submitted or transmitted to the International Bureau but not in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

0505/7052



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04380057.2

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr.:
Application no.: 04380057.2
Demande no:

Anmeldetag:
Date of filing: 12.03.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

ELI LILLY AND COMPANY
Lilly Corporate Center
Indianapolis, Indiana 46285
ETATS-UNIS D'AMERIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Opioid receptor antagonists

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G07D491/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

OPIOID RECEPTOR ANTAGONISTS


5 The present invention is in the field of medicinal chemistry. The invention relates specifically to compounds useful as opioid antagonists, methods of treatment, methods of using, and pharmaceutical compositions thereof.

Background

10 Three types of opioid receptors, mu, kappa, and delta opioid receptors are generally reported. Recent evidence points to the interactions between receptor dimer combinations of mu, kappa and/or delta receptors (called heterodimers) as also contributing to opioid activity. Opioid receptors and their normal regulation or lack thereof, has been implicated in disease states including irritable bowel syndrome, nausea, vomiting, pruritic dermatoses, depression, smoking and alcohol addiction, sexual dysfunction, stroke and trauma in animals. Therefore it is not surprising that the
15 ability to antagonistically bind opioid receptors has been shown to produce ameliorative, preventative and/or treatment effects in animals including humans afflicted with one or more of these disease states.


20 More recently, certain antagonists of the opioid receptors have been found to increase metabolic energy consumption, and reduction of weight in obese rats while maintaining muscle mass. These findings indicate that an effective opioid antagonist may be useful in preventing, treating and/or ameliorating the effect of obesity. Considering the percentage of the population that is obese in Western societies and the indirect costs associated with treating the effects and symptoms of obesity and Related Diseases, the importance of these findings cannot be overstated.

25 Though many opioid antagonists have been disclosed, the search continues for alternative and/or improved or more effective antagonists having an overall benefit to the patient with little or no major side effects. U.S Patent No. 4,891,379 disclosed phenylpiperidine opioid antagonists useful for the treatment of diabetes and obesity. In particular, U.S. patent 4,891,379 disclosed the compound LY 255582 represented by the
30 structure



U.S patent No. 6,140,352 discloses the compound of formula

Formula 1




(1)

wherein the variables X_1 , X_2 , X_3 , R_1 , R_3 , R_4 , R_5 and R_6 are as described therein, as agonists of the beta adrenergic receptor useful for the treatment of diabetes and obesity.

Regardless of these and other disclosures of compounds useful as opioid receptor antagonists, or useful for the treatment of obesity, and/or diabetes by other mechanisms, there remains an unmet medical need for a safe, effective and/or alternate treatment or prophylaxis of diseases associated with opioid receptors, particularly obesity and Related Diseases.

Summary of the Invention

The present invention provides a compound of the formula (I)



(I)

j is 1 or 2;

y is 0, 1, or 2; and z is 0, 1, or 2;

p is 0, 1, or 2;

wherein E is O or NH; and wherein each of

X_1 , X_2 , X_3 , X_4 , X_5 , or X_6 , is C, CH, or N; provided that each of rings A or C has no more than 2 nitrogen atoms; and provided that Ring B has 0 or 1 double bond;

R^1 and R^2 are independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, C_1 - C_{10} alkylaryl, $C(O)C_1$ - C_8 alkyl, SO_2C_1 - C_8 alkyl, SO_2C_1 - C_8 alkyl NR^8R^8 , $(CH_2)_nC(O)NR^8R^8$, SO_2C_1 - C_{10} alkylaryl, SO_2C_1 - C_8 alkylheterocyclic, C_4 - C_{10} alkylcycloalkane, $(CH_2)_nC(O)OR^8$, and $(CH_2)_nC(O)R^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C_1 - C_8 alkyl, C_2 - C_8 alkenyl, phenyl, C_3 - C_8 cycloalkyl, C_1 - C_8 alkylaryl, and $C(O)C_1$ - C_8 alkyl; and wherein R^1 and R^2 may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of oxo, amino, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, C_1 - C_8 alkylaryl, $C(O)C_1$ - C_8 alkyl, $CO(O)C_1$ - C_8 alkyl, halo, C_1 - C_8 haloalkyl;

R^3 and R^3 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, aryl, C_1 - C_8 alkylcycloalkyl, and C_1 - C_8 alkylaryl;

R^a and R^b are each independently selected from hydrogen, and C_1 - C_3 alkyl or combine with their respective carbon atoms to form the vinyl diradical $-CH=CH-$;

R^4 and R^5 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, C_1 - C_8 alkoxy, halo, C_1 - C_8 haloalkyl, phenyl, aryl, C_1 - C_8 alkylaryl, $(CH_2)_mNSO_2C_1$ - C_8 alkyl, $(CH_2)_mNSO_2$ phenyl, $(CH_2)_mNSO_2$ aryl, $-C(O)C_1$ - C_8 alkyl, and $-C(O)OC_1$ - C_8 alkyl; wherein each R^4 and R^5 is attached to its respective ring only at carbon atoms; wherein m is 1 or 2; and n is 1, 2, or 3;

R^6 and R^7 are each independently selected from hydrogen, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, $C(O)C_1$ - C_8 alkyl, SO_2C_1 - C_8 alkyl, SO_2C_1 - C_8 alkylaryl, SO_2C_1 - C_8 alkylheterocyclic, aryl, C_1 - C_8 alkylaryl, C_3 - C_7 cycloalkane, C_1 - C_6 alkylcycloalkane, $(CH_2)_nC(O)OR^8$, $(CH_2)_nC(O)R^8$, $(CH_2)_mC(O)NR^8R^8$, and $(CH_2)_mNSO_2R^8$; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C_1 - C_8 alkyl, C_2 - C_8 alkenyl, phenyl, and C_1 - C_8 alkylaryl; and wherein R^6 and R^7 may independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle may optionally have substituents selected from the group consisting of oxo, amino, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, and C_1 - C_8 alkylaryl;

R⁸ is independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, phenyl, benzyl, and C₅-C₈ alkylaryl; or a pharmaceutically acceptable salt, solvate, prodrug, tautomers, enantiomer, racemate, diastereomer, or mixture of diastereomers thereof.

5 The present invention also provides a method for the prevention, treatment and/or amelioration of the symptoms of obesity and Related Diseases comprising administering a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

10 The present invention also provides a pharmaceutical formulation comprising a compound of formula I in association with a carrier, diluent and/or excipient.

15 The present invention also relates to a method for the treatment and/or prophylaxis of obesity and Related Diseases including eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, depression, anxiety, epileptic seizure, hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, rheumatoid arthritis, stroke, hyperlipidemia, hypertriglycemia, hyperglycemia, hyperlipoproteinemia, substance abuse, drug overdose, compulsive behavior disorders (such as paw licking in dog), and addictive behaviors such as for example, gambling, and alcoholism, comprising administering a therapeutically effective amount of a compound
20 of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture of diastereomers thereof.

25 The present invention relates to a compound of formula (I) useful for the manufacture of a medicament for the treatment, prevention and/or amelioration of symptoms associated with obesity and Related Diseases.

30 In another embodiment, the present invention relates to a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture thereof, useful as an appetite suppressant.

35 The present invention relates to a method of achieving weight loss while maintaining lean muscle mass or minimizing the loss of lean muscle mass comprising administering a compound of formula I or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixture thereof, to a patient in need thereof.

The present invention provides a compound of formula I useful singly or in combination with other agents approved for the treatment, prevention and/or amelioration of obesity and related diseases and symptoms thereof.

Detailed Description of the Invention

As used herein, the term "patient" includes human and non-human animals such as companion animals (dogs and cats and the like) and livestock animals.

The preferred patient of treatment, amelioration and/or prevention of obesity and Related Diseases is a human.

The terms "treating" and "treat", as used herein, include their generally accepted meanings e.g. preventing, prohibiting, restraining, alleviating, ameliorating, slowing, stopping, or reversing the progression or severity of a pathological condition, or sequela thereof, described herein.

The terms "ameliorating" "preventing", "prevention of", "prophylaxis", "prophylactic" and "prevent" are used herein interchangeably and refer to reducing the severity of obesity and Related Diseases and the symptoms associated therewith, in a patient afflicted with same or reducing the likelihood that the recipient of a compound of formula I will be afflicted with or develop any of the pathological conditions or sequela thereof described herein.

As used herein, the term "effective amount" is synonymous with "effective dose" and means an amount of a compound of formula I that is sufficient in one or more administrations for preventing, ameliorating or treating a condition, or detrimental effects thereof, herein described, or an amount of a compound of formula I that is sufficient for antagonizing the opioid receptors to achieve the objectives of the invention.

The term "pharmaceutically acceptable" is used herein as an adjective and means substantially non-deleterious to the recipient patient.

The term "Active Ingredient" as used herein means a compound of formula I or a combination of a compounds of formula I or a combination of a compound of formula I and a co-antagonist of the opioid receptor or a combination a compound of formula I and other effective anti-obesity, weight loss or antidiabetic agent.

The term "formulation", as in pharmaceutical formulation, or "pharmaceutical composition" is intended to encompass a product comprising the Active Ingredient (as defined supra), and the inert ingredient(s) that make up the carrier, as well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical formulations of the present invention encompass any effective composition made by admixing a compound of the present invention and a pharmaceutical carrier. The pharmaceutical formulations of the present

invention also encompass a compound of the formula I and a pharmaceutically acceptable co-antagonist of opioid receptors useful for the treatment and/or prevention of obesity or Related Diseases.

5 The term "Related Diseases" as used herein refers to such symptoms, diseases or conditions caused by, exacerbated by, induced by or adjunct to the condition of being obese. Such diseases, conditions and/or symptoms include but are not limited to eating disorders (bulimia, anorexia nervosa, etc.), diabetes, diabetic complications, diabetic retinopathy, sexual/reproductive disorders, depression (particularly that induced by the awareness and loss of self esteem associated with obesity), anxiety, epileptic seizure, 10 hypertension, cerebral hemorrhage, congestive heart failure, sleeping disorders, atherosclerosis, rheumatoid arthritis, stroke, hyperlipidemia, hypertriglycemia, hyperglycemia, and hyperlipoproteinemia.

As used herein "other agents" approved for the treatment of obesity and /or related disease, or useful for weight loss and/or appetite suppression include but are not 15 limited to Xenical®, Meridia®, Lipitor®, Crestor®, Pravachol®, Zetia®, cannabinoid receptor antagonists, and other opioid receptor antagonists.

The term "suitable solvent" refers to any solvent, or mixture of solvents, inert to the ongoing reaction that sufficiently solubilizes the reactants to afford a medium within which to effect the desired reaction.

20 The term "mutual solvent" means a solvent that is used to dissolve sufficiently, two or more components of a reaction or mixture separately prior to reaction or mixing, that is a solvent common to more than one reagents or components of a mixture.

The term "nitrogen containing heterocycle" refers to a monocycle which is a 4, 5, 6, or 7-member ring containing 1, 2 or 3 nitrogen atoms in addition to the carbon atoms 25 completing the ring size, or a combination of 1 nitrogen atom and 1, or 2 atoms selected from oxygen, and sulfur in addition to the appropriate number of carbon atoms completing the ring size. A nitrogen containing heterocycle as used here may have 0, 1, 2 or 3 double bonds. A nitrogen containing heterocycle may be attached to or fused to an existing ring substituent thus forming a bicyclic or tricyclic ring system. Nonetheless, 30 the direct result of the formation of a nitrogen containing heterocycle by the joining of two groups and the nitrogen atom to which they are attached is to form a monocycle.

The term "C₁-C₈ alkyl" or C₁₋₈ alkyl" refers to and includes all groups, structural isomers and /or homologues of alkyl groups having from 1 to 8 carbon atoms. When the term C₁-C₈ alkyl precedes or prefixes another group, the term C₁-C₈ alkyl, only limits the 35 number of carbon atoms in the alkyl component. For example C₁-C₈ alkylaryl means an

aryl group having a C₁-C₈ alkyl group substituent such that the number of carbon atoms in the group C₁-C₈ alkylaryl is effectively the number of carbon atoms in the aryl group plus the number of carbon atoms in the C₁-C₈ alkyl group. Similarly, the term "C₁-C₈ alkylcycloalkyl" refers to a cycloalkane group having a C₁-C₈ alkyl substituent, and wherein the entire group C₁-C₈ alkylcycloalkane may itself be a substituent attached at either the alkyl group or the cycloalkyl group to a substrate. The definition and usage applies equally to other homologues of C₁-C₈ such as for example, C₁-C₇, C₁-C₆ etc.

The term "cycloalkane" or "cycloalkyl" means cycloalkanes having from 3 to 8 carbon atoms i.e. from cyclopropane to cyclooctane.

The term "hal" or "halo" as used herein refers to a halogen including fluorine, chlorine, bromine or iodine.

As used herein the terms "alkenyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon double bonds.

As used herein the terms "alkynyl" refers to straight or branched carbon atoms having 1 or 2 carbon-carbon triple bonds.

As used herein the term "alkoxy" refers to the group "O-alkyl" wherein alkyl is as defined previously.

The term "aryl" as used herein refers to compounds or groups having the Huckel 4n+2 pi electron arrangement and includes phenyl, benzyl, naphthyl, but excludes carbazoles and other fused tricyclic ring structures.

It is understood by one of skill in the art that where a substituent is absent, a hydrogen atom is indicated to achieve the required valency unless otherwise indicated. For example, if y is 0, then R⁴ is absent, and all applicable positions on the ring have hydrogen atoms to achieve the required valency for atoms in the ring.

As used herein, the term "protecting group" refers to a groups useful for masking reactive sites in a molecule to enhance the reactivity of another group or allow reaction at another desired site or sites following which the protecting group may be removed. Protecting groups are usually used to protect or mask groups including but not limited to -OH, -NH, and -COOH. Suitable protecting groups are known to one of skill in the art and are described in Protecting groups in Organic Synthesis, 3rd edition, Greene, T. W.; Wuts, P.G.M. Eds.; John Wiley and Sons, New York, 1999.

As used herein, the term "solvate" is a form of the compound of the invention wherein a crystal or crystals of a compound of the invention have been formed from a stoichiometric or non-stoichiometric amount of the compound of formula I and a solvent.

Typical solvating solvents include for example, water, methanol, ethanol, acetone and dimethylformamide.

In those instances where a compound of the invention possesses acidic or basic functional groups, various salts may be formed which are more water soluble and/or more physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, calcium, magnesium, aluminum and the like. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion-exchange resin.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," J. Phar. Sci., 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, hydrobromide, camsylate, carbonate, clavulanate, citrate, chloride, edetate, edisylate, estolate, esylate, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrochloride, hydroxynaphthoate, hydroiodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, sulfite, sulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pantothenate, phosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, tosylate, trifluoroacetate, trifluoromethane sulfonate, and valerate.

A compound of the invention as illustrated by formula I may occur as any one of its positional isomers, stereochemical isomers or regio-isomers, or canonical forms or tautomers, all of which are objects of the invention. Certain compounds of the invention may possess one or more chiral centers, and thus, may exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group, there exist the possibility of cis- and trans- isomeric forms of the compounds. The R- and S- isomers and mixtures thereof, including racemic mixtures as well as mixtures of enantiomers or cis- and trans- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art


10

20

15


For the groups R^1 and R^2

Preferred R^1 and R^2 groups are independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, and isopropyl. Also preferred are R^1 and R^2 groups independently selected from the group consisting of methyl, ethyl, propyl, isopropyl, phenyl,



each of which is optionally substituted with a group selected from the group consisting of halogen, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkyl heterocycle; or combine with a group selected from C₁-C₈ alkyl, halogen, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkyl heterocycle to form a substituted or unsubstituted bicycle or tricycle.

Also preferred are R¹ and R² groups which combine with each other to form a group selected from the group consisting of



each of which is optionally substituted with a group selected from the group consisting of halogen, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C₁-C₈ thioalkyl, C₁-C₈ alkylamino, phenyl, C₁-C₈ alkylsubstituted phenyl, C₄-C₈ heterocycle or C₁-C₄ alkylheterocycle.

Preferred R³ and R^{3'} Groups

A preferred R³ is hydrogen. A preferred R^{3'} group is selected from hydrogen, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, phenyl and benzyl. More preferably, both R³ and R^{3'} are hydrogen.

Preferred R⁴ Groups

A preferred R⁴ group is selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ haloalkyl, C₁-C₅ alkoxy, C₁-C₅ alkylamino, phenyl, C₁-C₅ alkylphenyl, C₁-C₅ alkylcycloalkyl, and C₁-C₅ thioalkyl. More preferred is a R⁴ group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. Most preferred is an R⁴ group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, propoxy, isopropoxy, and benzyl.

Though the groups R⁴ and a R⁵ may exist as multiple substituents on their respective ring substrates, a preferred embodiment of the invention involves compounds

wherein each of R^4 , and R^5 are independently absent, or singly substituted on their respective ring substrates.

Preferred R^a and R^b groups

Preferably R^a and R^b are each hydrogen, or combine to form the vinylic diradical $CH=CH$ when p is 2.

Preferred R^5 Groups

A preferred R^5 group is selected from the group consisting of hydrogen, halo, C_1 - C_5 alkyl, C_1 - C_5 haloalkyl, C_1 - C_5 alkoxy, C_1 - C_5 alkylamino, phenyl, C_1 - C_5 alkylphenyl, C_1 - C_5 alkylcycloalkyl, and C_1 - C_5 thioalkyl. More preferred is an R^5 group selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, chloro, fluoro, trifluoromethyl, methoxy, ethoxy, thiomethyl, phenyl, and benzyl. A most preferred R^5 group is selected from the group consisting of hydrogen, methyl, ethyl, isopropyl, fluoro, chloro, trifluoromethyl, methoxy, ethoxy, trifluoromethoxy, and benzyl.

Preferred R^6 and R^7 Groups

Preferred are R^6 and R^7 groups independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, pentyl, isopropyl, phenyl and benzyl.

Also preferred are compounds of formula I wherein R^6 and R^7 independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle optionally has substituents selected from the group consisting of oxo, amino, C_1 - C_8 alkyl, C_2 - C_8 alkenyl, C_2 - C_8 alkynyl, phenyl, C_1 - C_8 alkylaryl, $C(O)C_1$ - C_8 alkyl, $CO(O)C_1$ - C_8 alkyl, hydroxy, C_1 - C_8 alkoxy, halo, and haloalkyl.

Most preferred are compounds of the invention wherein R^6 and R^7 are both hydrogen.

Preferred E group

A most preferred E group is an oxygen atom (O).

Preferred A-ring

A preferred A-ring is a phenyl ring or a pyridine ring.

Preferred B ring

A preferred B ring is a 5, 6, or 7 membered ring. Also preferred is a B ring wherein $(CR^aR^b)_p$ equals $CH=CH$.

Preferred C-ring

A preferred C-ring is a phenyl ring, a pyrazine ring, a pyrimidine ring or a pyridine ring. Most preferred C ring is a phenyl ring or a pyridine ring.

Preferred values for n, m, and p

A preferred value for n is 1 or 2.

A preferred value for m is 1 or 2.

A preferred value for p is 2.

A preferred compound according to the present invention is a compound selected from the group consisting of:

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(Isobutylamino-methyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(4-Methyl-pentylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(2-Thiophen-2-yl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Pentylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Hexylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(Cyclohexylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Cyclooctylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Cycloheptylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(Cycloheptylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide trifluoroacetate salt,

8-[[2-(Tetrahydro-pyran-4-yl)-ethylamino]-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3,3-Dimethyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(2-Cyclopentyl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Morpholin-4-yl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Ethoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

5 8-[(2-Diethylamino-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Methoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide, and

10 8-[(3-Phenyl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(3-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(3-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

15 8-[2-(4-Chloro-phenyl)-pyrrolidin-1-ylmethyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

20 8-(2-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Phenyl-azepan-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Benzyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

25 8-[(3-Methyl-butylamino)-methyl]-dibenzofuran-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide,

30 8-[(3-Methyl-butylamino)-methyl]-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide,

or a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer and diastereomeric mixture thereof..


Preparing Compounds of the Invention

Compounds of formula I may be prepared as described in the following schemes and/or examples or following a combination of schemes known to one of skill in the art for making fragments and combinations thereof. Compounds employed as initial starting materials in the synthesis of compounds of the invention are well known and, to the extent not commercially available, are readily synthesized using specific references provided, or by standard procedures commonly employed by those of ordinary skill in the art and/or found in general reference texts.

More particularly, the compounds of the invention are produced in accordance with schemes 1 through 3 that are described in detail below, or analogous methods thereof. These reactions are often carried out following known procedures, methods, or analogous methods thereof. Examples of such known procedures and/or methods include those described in general reference texts such as Comprehensive Organic Transformations, VCH Publishers Inc, 1989; Compendium of Organic Synthetic Methods, Volumes 1-10, 1974-2002, Wiley Interscience; Advanced Organic Chemistry, Reactions Mechanisms, and Structure, 5th Edition, Michael B. Smith and Jerry March, Wiley Interscience, 2001; Advanced Organic Chemistry, 4th Edition, Part B, Reactions and Synthesis, Francis A. Carey and Richard J. Sundberg, Kluwer Academic / Plenum Publishers, 2000, etc., and references cited therein.

Compounds of the present invention are generally prepared starting with the corresponding dibenzo oxygenated tricyclic compound which is dibrominated to afford regioisomer 2, shown in scheme 1.


Scheme 1



Reaction with copper cyanide affords the dinitrile analog **3**. De-symmetrization is achieved by mono-oxidation to the carboxamide **4**. Preferably, a 50% but less than equivalent mole of reagents i.e. $\text{H}_2\text{O}_2/\text{K}_2\text{CO}_3$ is utilized to maximize de-symmetrization and minimize formation of the bis carboxamide compound. Further transformation of the remaining nitrile to an aldehyde with diisobutylaluminum hydride (DIBAL®) affords the compound **5**. The compound **5** upon reductive amination affords the desired product **6**. Details of specific procedures are provided in the scheme below and/or in the experimental section. The compound **6** may be converted to other compounds of the invention by elaborating the carboxamide functionality to N-substituted or N,N-disubstituted amide groups using procedures known to one of skill in the art and requiring minimal experimentation or research.

A preferred synthetic route for preparing compounds of formula I wherein Ring C is pyridine and the central ring (B) is a seven membered ring is shown in scheme 2.

Scheme 2




According to Scheme 2, 6-amino nicotinonitrile (**7**) is regioselectively brominated in the 5-position to afford the bromo compound **8**. The free amine functionality of compound **8** is converted to the chloride as in compound **9**. Compound **9** is coupled with the corresponding 3-bromo-4-hydroxybenzylic, N-tertbutyloxycarbonyl protected amine (**10**) to afford the coupled compound **11**. Palladium catalyzed cross coupling of **11** with vinyl tributyl stannane affords the arylvinyl compound **12** (see Stille Coupling, *Angew Chemie, Int. Ed.* 25, 508, (1986) and related references). The divinyl derivative **12** undergoes an intramolecular methathesis affording the compound **13** (see Grubbs second generation *Tetrahedron* 54, 4413, (1998) and related references therein). The compound **13** is hydrolyzed under basic conditions to afford the carboxamide **14**. For example, treatment of compound **13** with hydrogen peroxide in the presence of a base such as sodium carbonate in dimethylsulfoxide affords the carboxamide compound **14**. The carboxamide **14** is then de-protected at the amine by reaction with, for example, trifluoroacetic acid in methylene chloride preferably at room temperature to afford **15**, a compound of the invention. The carboxamide **14** is reduced under hydrogen atmosphere using palladium on carbon as catalyst to afford compound **16**. Compound **16** is then de-

protected at the amine by reaction with trifluoroacetic in methylene chloride to afford 17, a compound of the invention.

Alternatively, compound 17 is utilized to afford a disubstituted compound of the invention as shown in Scheme 3 below.


Scheme 3



As shown in Scheme 3, compound 17 upon reductive amination affords the amine 18 (wherein neither R¹ nor R² is hydrogen).


Substituted analogs of compound 1 may be prepared following for example, a scheme such as Scheme 4

Scheme 4



As shown in scheme 4, an optionally substituted Boc-protected amino 2-bromoaminophenol **19** may be coupled with an optionally substituted 3-bromo, 4-fluorobenzonitrile **20** to afford the coupled product **21**. The coupling reaction is accomplished under basic reaction conditions as described previously. The coupled product **21** is reacted with vinyl tributyl stannane under Stille coupling conditions described previously to afford the vinylic compound **22**. The compound **22** is then ring closed using the Grubbs methatesis procedure to afford the tricyclic compound **23**. The cyano group of the compound **23** is hydrolyzed under basic conditions discussed previously to afford the carboxamide compound **24**. The carboxamide compound **24** is reduced at the alkenyl group to afford the reduced compound **25**, which is further deprotected at the boc group to afford the compound **27**. Alternatively, the carboxamide **24** is hdeprotected at eh amino group to afford the alekeny compound **26**. One of skill in the art is aware that the use of a disubstitued amine compound as staarting material in place of compound **19** would result following the protecool above in the disubstituted amine analog of compound **26** or **27**.

An example of a compound of formula **20** is 3-bromo-3,4,5-trifluorobenzonitrile disclosed in European patent application EP 85/114373. Compoundsof formula **19** may be prepared by methods known to one of skill in the art or may be available from commercial sources. For example, compounds of formula **19** may be prepared as shown in scheme 6.



As shown in Scheme 6, a compound of formula **19** may be prepared starting with an optionally substituted hydroxybenzaldehyde **28**, which is reductively aminated to the amine **29**. The amine **29** is boc protected to afford the resulting amine **19**. Specific

examples of the optionally substituted compound 28 include compounds 30, 31, and 32 all of which are available from commercial sources.

Method of Using the Invention

5 As noted above, the compounds of the present invention are useful in blocking the effect of agonists at mu, kappa, and/or delta opioid receptors. As such, the present invention also provides a method for blocking a mu, kappa, delta receptor or receptor combination (heterodimer) thereof in a mammal comprising administering to said mammal a receptor blocking dose of a compound of formula I.

10 The term "receptor blocking dose", as used herein, means an amount of a compound of formula I necessary to block a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof following administration to a mammal requiring blocking of a mu, kappa, or delta receptor or receptor combination (heterodimer) thereof.

15 The compounds of formula I or combinations thereof, are effective over a wide dosage range. For example, dosages per day will normally fall within the range of about 0.05 to about 250 mg/kg of body weight. In the treatment of adult humans, the range of about 0.5 to about 100 mg/kg, in single or divided doses, is preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician in light of the relevant circumstances, including the condition to be treated, the choice of compound to be administered, the age, weight, response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration. Therefore, the above dosage ranges are not intended to limit the scope of the invention in any way. The compounds may be administered by a variety of routes such as the oral, transdermal, subcutaneous, intranasal, intramuscular and intravenous routes.

25 A variety of physiologic functions have been shown to be subject to or influenced by mu, kappa, or delta receptors or receptor combination (heterodimers) in the brain. As such, the compounds of the present invention are believed to have the ability to treat disorders associated with these receptors or combinations thereof such as eating disorders, opioid overdose, depression, smoking, alcoholism, sexual dysfunction, shock, stroke, spinal damage and head trauma. As such, the present invention also provides methods of treating the above disorders by blocking the effect of agonists at a mu, kappa, delta receptors or receptor combinations (heterodimer) thereof.

Assay Methodology

The compounds of the present invention have been found to display significant activity in an opioid receptor binding assay which measures the ability of the compounds to block the mu, kappa, delta or receptor combination (heterodimer) thereof. The functional antagonist potency (Kb) of the sample compounds was determined using the GTP γ S binding assay. GTP γ S – based functional assays provide an in vitro measure of the activity of opioid agonists and antagonists. Opioid reference compounds or test compound are incubated with membrane homogenate from cells expressing the cloned human mu, kappa or delta opioid receptor and radiolabeled [35S]GTP γ S. If the compound activates the opioid receptor, an increase in the binding of radiolabeled GTP γ S is observed. Similarly, if the compound exhibits antagonist activity, it interferes with the ability of the control agonist to stimulate GTP γ S binding. These tests provide an in vitro measurement of the activity of the compound at human opioid receptors.

GTP- γ -S Binding Assay

An SPA - based GTP- γ -S assay format was developed based on previous opioid (Emmerson et al., J. Pharm. Exp Ther 278, 1121, 1996; Horng et al., Society for Neuroscience Abstracts, 434.6, 2000) and muscarinic (DeLapp et al., JPET 289, 946, 1999) assay formats. Membranes were resuspended in 20 mM HEPES, 100 mM NaCl, 5 mM MgCl₂, 1 mM DTT, and 1 mM EDTA. Fifty (50) μ L of GTP- γ -[35S], compound, membrane suspension (20 microgram/well), and wheat germ agglutinin coated SPA beads (1mg/well) were added to clear bottom 96 well assay plates. GDP (200 mM) was added to the membrane solution prior to addition to the assay plates. Plates were sealed and incubated for four hours at room temperature then placed in a refrigerator overnight to allow the beads to settle. Signal stability at 4 °C was determined to be > 60 hours. Plates were warmed to room temperature and counted in a Wallac Microbeta scintillation counter. For antagonist assays, specific agonists were added at the following concentrations: (MOR) DAMGO 1 micromolar, (DOR) DPDPE 30 nM, (KOR) U69593 300 nM. Kb's were determined by Cheng-Prusoff equation (see Cheng and Prusoff, Biochem. Pharmacol. 22, 3099, 1973). Results obtained for a sample of compounds of the invention in the GTP- γ -S Binding Assay are shown in table 1 below.

Table 1

Example #	Kb (nM) mu	Kb (nM) kappa	Kb (nM) delta
1	1.2	6.1	11.0

2	12.0	29.8	
3	0.8	6.5	7.4
4	0.9	12.6	7.1
5	1.7	4.9	
6	0.7	4.4	10.3
12	0.7	2.2	2.1
13	4.0	12.1	14.4
14	0.6	8.6	7.0
17	2.1	25.7	26.7

Formulation

A compound of the invention is preferably presented in the form of a pharmaceutical formulation comprising a pharmaceutically acceptable carrier, diluent or excipient and a compound of the invention. Such compositions will contain from about 0.1 percent by weight to about 90.0 percent by weight of the compound of the invention (Active Ingredient). As such, the present invention also provides pharmaceutical formulations comprising a compound of the invention and a pharmaceutically acceptable carrier, diluent or excipient therefore.

In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material that acts as a vehicle, excipient or medium for the active ingredient. Thus, the composition can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, emulsions, solutions, syrups, suspensions, aerosols (as a solid or in a liquid medium), and soft and hard gelatin capsules.

Examples of suitable carriers, excipients, and diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, tragacanth, gelatin, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, water, and mineral oil. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or

delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

For oral administration, the Active Ingredient, a compound of this invention, may be admixed with carriers and diluents and molded into tablets or enclosed in gelatin capsules.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 1 to about 500 mg, more usually about 5 to about 300 mg, of the Active Ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier.

In order to more fully illustrate the operation of this invention, the following formulation examples are provided. The examples are illustrative only, and are not intended to limit the scope of the invention. The formulations may employ as Active Ingredient any of the compounds of the present invention.

FORMULATION 1

Hard gelatin capsules are prepared using the following ingredients:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	250	55
Starch dried	200	43
Magnesium stearate	10	2

The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities.

FORMULATION 2

Capsules each containing 20 mg of medicament are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	20	10
Starch	89	44.5
Microcrystalline cellulose	89	44.5
Magnesium stearate	2	1

The active ingredient, cellulose, starch and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve and filled into a hard gelatin capsule.

FORMULATION 3

Capsules each containing 100 mg of active ingredient are made as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	100	30
Polyoxyethylene Sorbitan monooleate	50mcg	0.02
Starch powder	250	69.98

The above ingredients are thoroughly mixed and placed in an empty gelatin capsule.

FORMULATION 4

Tablets each containing 10 mg of active ingredient are prepared as follows:

Compound	Amount per capsule (mg)	Concentration by weight (%)
Active Ingredient	10	10
Starch	45	45
Microcrystalline cellulose	35	35
Polyvinylpyrrolidone (as 10% solution in water)	4	4
Sodium carboxymethyl starch	4.5	4.5
Magnesium stearate	0.5	0.5
talc	1	1

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders, which are then passed through a No. 14 mesh U.S. sieve. The granule so produced is dried at 50-60 °C and passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules, which after mixing, is compressed on a tablet machine to yield a tablet weighing 100 mg.

FORMULATION 5

A tablet formula may be prepared using the ingredients below:

Compound	Amount per capsule (mg)	Percent by weight (%)
Active Ingredient	250	38
Cellulose microcrystalline	400	60
Silicon dioxide fumed	10	1.5
Stearic acid	5	0.5

The components are blended and compressed to form tablets each weighing 665mg.

FORMULATION 6

Suspensions each containing 5 mg of medicament per 5 ml dose are made as follows:

Compound	Amount per 5mL suspension (ml)
Active Ingredient	5
Sodium carboxymethyl cellulose	50
Syrup	1.25
Benzoic acid solution	0.10
Flavor	q.v.
Color	q.v.
Water	q.s. to 5mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethylcellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color is diluted with some of the water and added to the paste with stirring. Sufficient water is then added to produce the required volume.

FORMULATION 7

An aerosol solution is prepared containing the following components:

Compound	Concentration by weight (percent)
Active Ingredient	0.25
Ethanol	29.75
Propellant 22 (chlorodifluoromethane)	70.0

The active compound is mixed with ethanol and the mixture added to a portion of the Propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted further with the remaining amount of propellant. The valve units are then fitted to the container.

Examples


Compounds of the invention may be prepared following procedures disclosed herein or known modifications thereof. Unless otherwise indicated, reagents are generally available from chemical distributors including those specializing in fine limited use chemicals.

Example 1

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[*b,f*]oxepine-2-carboxilic acid amide

Step 1

2,8-Dibromo-10,11-dihydro-dibenzo[*b,f*]oxepine



To a solution of (10,11-dihydro-dibenzo[*b,f*]oxepine* (2.7 g, 13.8 mmol) in AcOH (11 mL) add Br₂ 2.0 M (17.2 mL, 13.8 mmol) in AcOH. Stir the mixture at 55°C for 4 hours. Cool the mixture at room temperature and eliminate the solvent under reduced pressure. Dissolve the crude in CH₂Cl₂ and wash with aqueous solution of NaHSO₃. Dry the


organic layer over Na_2SO_4 . Eliminate the solvent to obtain 4.9 g of the title compound (99%) which is used in the next step without purification.

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 7.29-7.23 (m, 4H), 7.04-6.99 (m, 2H), 3.08 (s, 4H). $^{13}\text{C-NMR}$ (CDCl_3 , 500 MHz): 155.9, 133.8, 133.2, 130.6, 122.9, 116.8, 30.9.

*The synthesis of (10,11-dihydro-dibenzo[*b,f*]oxepine is described in *J. Am. Chem. Soc.* 1969, 91, 1665.

Step 2

10,11-dihydro-dibenzo[*b,f*]oxepine-2,8-dicarbonitrile




To a solution of the intermediate above (4.9 g, 13.8 mmol) in dry DMF (120 mL) add CuCN (8.6 g, 96.8 mmol), heat the mixture at 160°C under N_2 atmosphere overnight. Cool the mixture at room temperature, add NH_4OH and bubble air during 4 hours. Add cold ($0-25^\circ\text{C}$) water and extract with EtOAc , dry over Na_2SO_4 . Eliminate the solvent under reduced pressure. Purify by flash chromatography on silica gel (eluent: hexane/ EtOAc 4/1) to afford the title compound (2.0 g, 59%).

$^1\text{H-NMR}$ (CDCl_3 , 300 MHz): 7.54-7.48 (m, 4H), 7.28-7.22 (m, 2H), 3.17 (s, 4H). $^{13}\text{C-NMR}$ (CDCl_3 , 300 MHz): 158.0, 134.7, 132.5, 131.8, 122.2, 118.5, 107.5, 31.1.

Step 3

8-Cyano-10,11-dihydro-dibenzo[*b,f*]oxepine-2-carboxylic acid amide




To a solution of the intermediate above (2.0 g, 8.1 mol) in dry DMSO (80 mL) add K_2CO_3 (562 mg, 4.06 mmol), cool the mixture at 0°C and add H_2O_2 33% (837 μL , 8.1 mmol) dropwise. Stir the mixture at room temperature overnight. Add cold ($0-25^\circ\text{C}$) water and extract with EtOAc , dry the organic layer over Na_2SO_4 , and eliminate the solvent at reduced pressure. Purify the crude product by flash column chromatography on silica gel (eluent: EtOAc) to obtain the title compound (1.08 g, 48%).

$^1\text{H-NMR}$ (MeOD, 300 MHz): 7.75-7.70 (m, 2H), 7.60-7.54 (m, 2H), 7.33-7.22 (m, 2H), 3.28 (m, 4H). $^{13}\text{C-NMR}$ (MeOD, 300 MHz): 167.4, 159.4, 157.7, 135.6, 133.1, 132.2, 131.2, 130.7 (2C), 127.5, 122.5, 120.9, 118.9, 106.9, 30.8, 30.6.

Step 4


8-Formyl-10,11-dihydro-dibenzo[*b,f*]oxepine-2-carboxylic acid amide



To a solution of the intermediate from step 3 (2.5g, 9.5 mmol) in dry CH_2Cl_2 (320 mL), cooled at -78°C and under N_2 atmosphere, add DIBAL 1.0M in toluene (28.4 mmol). Allow the mixture to warm to room temperature and stir overnight. Cool the mixture to 0°C and add $\text{AcOH}/\text{H}_2\text{O}$. Stir at room temperature for 1 hour. Extract the aqueous layer with CH_2Cl_2 (x2). Combine the organic layers and dry over Na_2SO_4 . Eliminate the solvent and purify by flash chromatography (eluent: EtOAc/hexane 3/1) to obtain the title compound (1.75 g, 69%). $^1\text{H-NMR}$ (MeOD, 300 MHz): 9.91 (s, 1H), 7.81-7.73 (m, 4H), 7.37-7.26 (m, 2H), 3.21 (m, 4). $^{13}\text{C-NMR}$ (MeOD, 300 MHz): 193.2, 162.6, 160.3, 134.4, 133.8, 133.2, 131.9, 131.3, 130.6, 128.8, 123.3, 122.4, 32.9, 32.8.

Step 5

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[*b,f*]oxepine-2-carboxylic acid amide



Combine intermediate step 4 (250 mg, 0.94 mmol) with 3-methylbutylamine (110 μ L, 0.94 mmol), MeOH (12 mL) and molecular sieves 3A (1.5 g). Stir at room temperature overnight. Add NaBH₄ (178 mg, 4.7 mmol) and stir at room temperature for 2 hours. Filtrate over celite with MeOH and eliminate the solvent. Purify using SCX ion-exchange chromatography to afford the title compound (300 mg, 91%).

By the method of example 1, using the corresponding reagents, the following compounds (examples 2-20) were prepared. The purification process depends on the structure of the corresponding example.

Example	Name	Electronspray MS M+1 ion	¹ H-NMR (300 MHz, CD ₃ OD)
1	8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	339	7.70-7.66 (m, 2H), 7.20-7.08 (m, 4H), 3.70 (s, 2H), 3.14 (m, 4H), 2.63-2.55 (m, 2H), 1.63-1.36 (m, 3H), 0.89 (d, 6H, J= 6.9 Hz)
2	8-(Isobutylamino-methyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	325	7.73-7.69 (m, 2H), 7.22-7.12 (m, 4H), 3.75 (s, 2H), 3.17 (bs, 4H), 2.44 (d, 2H, J= 6.8 Hz), 1.91-1.78 (m, 1H), 0.93 (d, 6H, J= 6.7 Hz).
3	8-[(4-Methyl-pentylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	353	7.75-7.70 (m, 2H), 7.26-7.18 (m, 4H), 3.91 (s, 2H), 3.19 (bs, 4H), 2.79-2.74 (m, 2H), 1.67-1.53 (m, 2H), 1.31-1.20 (m, 3H), 0.92 (d, 6H, J= 6.7 Hz)
4	8-[(2-Thiophen-2-yl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	379	7.64-7.59 (m, 2H), 7.12-7.04 (m, 5H), 6.85-6.82 (m, 1H), 6.76 (m, 1H), 3.64 (s, 2H), 3.10-3.03 (m, 4H), 2.95 (t, 2H, J= 7.6 Hz), 2.77 (t, 2H, J= 7.3 Hz)
5	8-Pentylaminomethyl-	339	7.64-7.59 (m, 2H), 7.12-7.02


	10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide		(m, 4H), 3.61 (s, 2H), 3.07 (bs, 4H), 2.50-2.45 (m, 2H), 1.49-1.40 (m, 2H), 1.30-1.15 (m, 4H), 0.83 (t, 3H, J= 6.9 Hz)
6	8-Hexylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	353	7.64-7.59 (m, 2H), 7.12-7.02 (m, 4H), 3.62 (s, 2H), 3.07 (bs, 4H), 2.51-2.46 (m, 2H), 1.47-1.39 (m, 2H), 1.28-1.14 (m, 6H), 0.80 (t, 3H, J= 6.9 Hz)
7	8-[(Cyclohexylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	365	7.63-7.59 (m, 2H), 7.12-7.01 (m, 4H), 3.59 (s, 2H), 3.07 (bs, 4H), 2.30 (d, 2H, J= 6.8 Hz), 1.69-1.57 (m, 5H), 1.46-1.35 (m, 1H), 1.27-1.09 (m, 3H), 0.87-0.75 (m, 2H)
8	8-Cyclooctylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	379	7.63-7.59 (m, 2H), 7.12-7.01 (m, 4H), 3.60 (s, 2H), 3.07 (bs, 4H), 2.64-2.58 (m, 1H), 1.76-1.31 (m, 14H).
9	8-Cycloheptylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	365	7.63-7.59 (m, 2H), 7.12-7.01 (m, 4H), 3.60 (s, 2H), 3.07 (bs, 4H), 2.60-2.51 (m, 1H), 1.86-1.78 (m, 2H), 1.65-1.31 (m, 10H)
10	8-[(Cycloheptylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	379	7.64-7.59 (m, 2H), 7.12-7.01 (m, 4H), 3.59 (s, 2H), 3.07 (bs, 4H), 2.30 (d, 2H, J= 6.5 Hz), 1.68-1.04 (m, 13H)

	trifluoroacetate salt		
12	8-[[2-(Tetrahydro-pyran-4-yl)-ethylamino]-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	381	7.75-7.71 (m, 2H), 7.24-7.13 (m, 4H), 3.41 (dd, 2H, J= 4.0 and 11.3 Hz), 3.73 (s, 2H), 3.40 (dt, 2H, J= 1.6 and 11.7 Hz), 3.18 (bs, 4H), 2.66-2.61 (m, 2H), 1.63-1.46 (m, 5H), 1.34-1.25 (m, 2H)
13	8-[(3,3-Dimethyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	353	7.64-7.59 (m, 2H), 7.12-7.02 (m, 4H), 3.61 (s, 2H), 3.06 (bs, 4H), 2.54-2.49 (m, 2H), 1.39-1.34 (m, 2H), 0.92 (s, 9H)
14	8-[(2-Cyclopentyl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	365	7.64-7.59 (m, 2H), 7.13-7.02 (m, 4H), 3.62 (s, 2H), 3.07 (bs, 4H), 2.53-2.48 (m, 2H), 1.76-1.44 (m, 9H), 1.07-1.00 (m, 2H)
15	8-[(2-Morpholin-4-yl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	382	7.64-7.59 (m, 2H), 7.13-7.02 (m, 4H), 3.63 (s, 2H), 3.59-3.55 (m, 4H), 3.07 (bs, 4H), 2.60 (t, 2H, J= 6.9 Hz), 2.41 (t, 2H, J= 6.6 Hz), 2.34-2.31 (m, 4H)
16	8-[(3-Morpholin-4-yl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	396	7.64-7.59 (m, 2H), 7.12-7.02 (m, 4H), 3.61 (s, 2H), 3.59-3.55 (m, 4H), 3.07 (bs, 4H), 2.54 (t, 2H, J= 7.2 Hz), 2.37-2.27 (m, 6H), 1.68-1.58 (m, 2H)

17	8-[(3-Ethoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	355	7.62-7.58 (m, 2H), 7.10-6.99 (m, 4H), 3.59 (s, 2H), 3.39-3.32 (m, 4H), 3.02 (bs, 4H), 2.57 (t, 2H, J= 6.9 Hz), 1.72-1.63 (m, 2H), 1.05 (t, 3H, J= 7.2 Hz)
18	8-[(2-Diethylamino-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	368	7.64-7.59 (m, 2H), 7.13-7.03 (m, 4H), 3.64 (s, 2H), 3.07 (bs, 4H), 2.62-2.43 (m, 8H), 0.94 (t, 6H, J= 7.2 Hz)
19	8-[(3-Methoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	341	7.73-7.69 (m, 2H), 7.22-7.11 (m, 4H), 3.71 (s, 2H), 3.49 (t, 2H, J= 6.9 Hz), 3.33 (s, 3H), 3.19 (bs, 4H), 2.66 (t, 2H, J= 7.1 Hz), 1.83-1.74 (m, 2H)
20	8-[(3-Phenyl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	387	7.72-7.69 (m, 2H), 7.24-7.05 (m, 9H), 3.65 (s, 2H), 3.11 (bs, 4H), 2.63-2.54 (m, 4H), 1.87-1.77 (m, 2H)

Example 21

8-[[Cyclopropylmethyl-(3-methyl-butyl)-amino]-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide




To a solution of amine of example 1 (30 mg, 0.09 mmol) in MeOH-AcOH 5% (0.5 mL) add cyclopropanecarboxaldehyde (0.18 mmol). Stir the mixture at room temperature overnight. Add NaBH₃CN and stir the mixture at room temperature for about two hours. Add more aldehyde (1.5 mmol) and stir overnight. Purify by SCX ion-exchange chromatography to obtain the title compound (22 mg). ¹H-NMR (CD₃OD, 300 MHz): 7.72-7.69 (m, 2H), 7.21-7.09 (m, 4H), 3.64 (s, 2H), 3.14 (bs, 4H), 2.61-2.56 (m, 2H), 2.37 (d, 2H, J= 6.5 Hz), 1.60-1.37 (m, 4H), 0.87 (d, 6H, J= 6.7 Hz), 0.55-0.49 (m, 2H), 0.13-0.08 (m, 2H). Electrospray MS M+1 ion= 393.

By the method of example 21, using the corresponding reagents, the following compounds (examples 22-23) were prepared. The purification process depends on the structure of the corresponding example.

Example 22


8-[[Cyclohexylmethyl-(3-methyl-butyl)-amino]-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide



¹H-NMR (CD₃OD, 300 MHz): 7.72-7.69 (m, 2H), 7.21-7.07 (m, 4H), 3.44 (s, 2H), 3.16-3.12 (m, 4H), 2.61-2.56 (m, 2H), 2.36 (t, 2H, J= 7.1 Hz), 2.16 (d, 2H, J= 7.1 Hz), 1.84-1.13 (m, 12H), 0.83 (d, 6H, J= 6.8 Hz). Electrospray MS M+1 ion= 435.

Example 23


8-[[Methyl-(3-methyl-butyl)-amino]-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide trifluoroacetate salt



¹H-NMR (CD₃OD, 300 MHz): 7.76-7.72 (m, 2H), 7.37-7.23 (m, 4H), 4.43-4.16 (m, 2H), 3.21-3.10 (m, 6H), 2.79 (s, 3H), 1.67 (m, 3H), 0.98 (d, 6H; J= 6.7 Hz). Electrospray MS M+1 ion= 353.

Example 24

8-(3-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide



To a solution of aldehyde intermediate obtained in step 4, Example 1 (92 mg, 0.34 mmol) in THF/AcOH (6.5 mL/30 μ L) add 3-phenylpyrrolidine (0.34 mmol). Stir the mixture at room temperature overnight. Add NaBH(OAc)₃ (0.52 mmol) and stir at room temperature overnight. Eliminate the solvent and purify by ISCO chromatography (eluent: CHCl₃/B 0-10%; B: EtOH/NH₄OH 10%).

By the method of example 24, using the corresponding reagents, the following compounds (examples 25-30) were prepared. The purification process depends on the structure of the corresponding example.


Examp le	Name	Electronsp ray MS M+1 ion	¹ H-NMR (300 MHz, CDCl ₃)
24	8-(3-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	399	7.65 (d, 1H, J= 2.4 Hz), 7.57(dd, 1H, J= 2.4 and 8.5 Hz), 7.33-7.09 (m, 9H), 3.67 (s, 2H), 3.49-3.35 (m, 1H), 3.21-3.07 (m, 5H), 2.93-2.90 (m, 1H), 2.76-2.72 (m, 1H), 2.56-2.50 (m, 1H), 2.42-2.29 (m, 1H), 2.02-1.88 (m, 1H)
25	8-(3-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-	413	7.65 (d, 1H, J= 2.4 Hz), 7.56 (dd, 1H, J= 2.4 and 8.5 Hz),

	dibenzo[b,f]oxepine-2-carboxylic acid amide		7.30-7.08 (m, 9H), 3.49 (s, 2H), 3.19-3.11 (m, 4H), 3.00-2.83 (m, 3H), 2.07-1.90 (m, 3H), 1.76 (bs, 2H), 1.52-1.38 (m, 1H)
26	8-[2-(4-Chloro-phenyl)-pyrrolidin-1-ylmethyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	433	7.74-7.70 (m, 2H), 7.43 (d, 2H, J= 8.5 Hz), 7.33 (d, 2H, J= 8.6 Hz), 7.20 (d, 1H, J= 8.3 Hz), 7.08 (bs, 3H), 3.68 (d, 1H, J= 12.9 Hz), 3.40-3.33 (m, 1H), 3.18-3.05 (m, 6H), 2.31-2.19 (m, 2H), 1.92-1.62 (m, 3H)
27	8-(2-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	399	7.65 (d, 1H, J= 2.0 Hz), 7.56 (dd, 1H, J= 2.4 and 8.5 Hz), 7.46-7.43 (m, 2H), 7.36-7.31 (m, 2H), 7.27-7.17 (m, 2H), 7.07-7.03 (m, 3H), 3.76 (d, 1H, J= 12.9 Hz), 3.33 (t, 1H, J= 8.5 Hz), 3.16-3.05 (m, 4H), 2.98 (d, 1H, J= 12.9 Hz), 2.24-2.12 (m, 2H), 1.95-1.66 (m, 4H)
28	8-(2-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	413	7.56 (d, 1H, J= 2.0 Hz), 7.47 (dd, 1H, J= 2.4 and 8.1 Hz), 7.36-7.33 (m, 2H), 7.23 (t, 2H, J= 6.9 Hz), 7.16-7.08 (m, 2H), 6.97-6.90 (m, 3H), 3.59 (d, 1H, J= 13.3 Hz), 3.08-2.96 (m, 5H), 2.88-2.84 (m, 1H), 2.66 (d, 1H, J= 13.3 Hz), 1.82 (td, 1H, J= 2.7 and 11.3 Hz), 1.70-0.74 (m, 6H)
29	8-(2-Phenyl-azepan-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-	427	7.56 (d, 1H, J= 2.0 Hz), 7.47 (dd, 1H, J= 2.0 and 8.1 Hz), 7.39-7.36 (m, 2H), 7.22 (t, 2H,

	carboxylic acid amide		J= 6.9 Hz), 7.13-7.08 (m, 2H), 7.04-6.93 (m, 3H), 3.60 (t, 1H, J= 6.1 Hz), 3.49 (d, 1H, J= 14.1 Hz), 3.23 (d, 1H, J= 13.7 Hz), 3.08-2.98 (m, 4H), 2.79 (dd, 1H, J= 6.9 and 14.5 Hz), 2.59 (dd, 1H, J= 1H, J= 6.9 and 14.1 Hz), 1.86-1.40 (m, 8H)
30	8-(2-Benzyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide	413	7.57 (d, 1H, J= 2.4 Hz), 7.48 (dd, 1H, J= 2.4 and 8.5 Hz), 7.21-7.00 (m, 9H), 3.95 (d, 1H, J= 12.9 Hz), 3.15-2.94 (m, 5H), 2.84 (dt, 1H, J= 2.4 and 8.3 Hz), 2.60-2.41 (m, 2H), 2.06 (q, 1H, J= 8.1 Hz), 1.71-1.44 (m, 5H)

Example 31


8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxilic acid amide hydrochloride salt



Dissolve amine of example 1 in 1N hydrochloric acid in AcOEt. Isolate the white precipitate by vacuum filtration to obtain the title compound. ¹H-NMR (CD₃OD, 300 MHz): 7.91-7.86 (m, 2H), 7.50-7.37 (m, 4H), 4.31 (s, 2H), 3.36 (s, 4H), 3.24-3.18 (m, 2H), 1.89-1.70 (m, 3H), 1.13 (d, 6H, J= 6.5 Hz). Electrospray MS M+1 ion= 339.

Example 32


8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxilic acid amide methanesulfonate salt



Dissolve amine of example 1 in THF (0.1M solution). Add metanesulfonic acid (1 equiv). Isolate the white precipitate by vacuum filtration to obtain the title compound. ¹H-NMR (CD₃OD, 300 MHz): 7.90-7.87 (m, 2H), 7.51-7.37 (m, 4H), 4.31 (s, 2H), 3.36 (s, 4H), 3.22-3.19 (m, 2H), 2.87 (s, 3H), 1.86-1.73 (m, 3H), 1.14 (d, 6H, J= 6.5 Hz). Electrospray MS M+1 ion= 339.

Example 33


8-[(2-Morpholin-4-yl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide dimethanesulfonate salt



Dissolve amine of example 15 in THF (0.1M solution). Add metanesulfonic acid (1 equiv). Isolate the white precipitate by vacuum filtration. ¹H-NMR (CD₃OD, 300 MHz): 7.73-7.69 (m, 2H), 7.42-7.38 (m, 2H), 7.27-7.20 (m, 2H), 4.27 (s, 2H), 4.04 (bs, 2H), 3.88 (bs, 2H), 3.58 (m, 6H), 3.18 (m, 6H), 2.70 (s, 6H). Electrospray MS M+1 ion= 382.

Example 34

8-[(3-Morpholin-4-yl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide dimethanesulfonate salt




Dissolve amine of example 16 in THF (0.1M solution). Add metanesulfonic acid (1 equiv). Isolate the white precipitate by vacuum filtration. ¹H-NMR (CD₃OD, 300 MHz): 7.73-7.69 (m, 2H), 7.38-7.35 (m, 2H), 7.27-7.20 (m, 2H), 4.20 (s, 2H), 3.86 (bs, 4H), 3.19-3.13 (m, 12H), 2.71 (s, 6H), 2.22-2.06 (m, 2H). Electrospray MS M+1 ion= 396.

Example 35

8-[(3-Methyl-butylamino)-methyl]-dibenzofuran-2-carboxylic acid amide

Step 1

Dibenzofuran-2,8-dicarbonitrile




To a solution of 2,8-Diiodo-dibenzofuran (763 mg, 1.8 mmol) in dry DMF (16 mL) add CuCN (1.1 g, 12.7 mmol), heat the mixture at 160°C under N₂ atmosphere overnight. Cool the mixture at room temperature, add NH₄OH and bubble air during 4 hours. Add cool water and extract with EtOAc, dry over Na₂SO₄. Eliminate the solvent under reduced pressure. Purify by flash chromatography on silica gel (eluent: hexane/EtOAc 3/1) to afford the title compound (25 mg, 6%).

¹H-NMR (DMSO, 200 MHz): 8.81 (dd, 2H, J= 0.8 and 1.6 Hz), 8.11-8.06 (m, 2H), 8.03-7.99 (m, 2H).

Step 2

8-Cyano-dibenzofuran-2-carboxylic acid amide




To a solution of the intermediate above (22 mg, 0.1 mmol) in dry DMSO (1 mL) add K₂CO₃ (7 mg, 0.05 mmol), cool the mixture at 0°C and add H₂O₂ 33% (11 µl, 0.1 mmol)

dropwise. Stir the mixture at room temperature overnight. Add cool water and extract with EtOAc, dry the organic layer over Na₂SO₄, eliminate the solvent at reduced pressure. Purify by flash chromatography on silica gel (eluent: EtOAc) to obtain the title compound (15 mg, 62%).

¹H-NMR (DMSO, 300 MHz): 8.79 (d, 2H, J= 1.2 Hz), 8.13 (dd, 1H, J= 1.8 and 8.7 Hz), 8.11 (bs, 1H), 8.03 (dd, 1H; J= 1.4 and 8.5 Hz), 7.98 (d, 1H, J= 8.3 Hz), 7.86 (d, 1H, J= 8.9 Hz), 7.49 (bs, 1H).

Step 3


8-Formyl-dibenzofuran-2-carboxylic acid amide



To a suspension of the intermediate above (15 mg, 0.07 mmol) in dry CH₂Cl₂ (3 mL), cooled at -78°C and under N₂ atmosphere, add DIBAL 1.0M in toluene (0.25 mmol). Stir the mixture for 1 hour. Add AcOH/H₂O. Stir at room temperature for 1 hour. Extract the aqueous layer with EtOAc (x2), combined organic layers and dry over Na₂SO₄. Eliminate the solvent and purify by flash chromatography (eluent: EtOAc/hexane 4/1) to obtain the title compound (10 mg, 61%). ¹H-NMR (MeOD, 300 MHz): 10.15 (s, 1H), 8.73 (m, 2H), 8.18-8.13 (m, 2H), 7.84 (d, 1H, J= 8.7 Hz), 7.76 (dd, 1H, J= 0.6 and 8.7 Hz).

Step 4

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide



Combine intermediate above (10 mg, 0.04 mmol) with 3-methylbutylamine (5 μL, 0.04 mmol), MeOH (1 mL) and molecular sieves 3A (60 mg) and stir at room temperature

overnight. Add NaBH_4 (8 mg, 0.2 mmol) and stir at room temperature for 2 hours. Filtrate over celite with MeOH and eliminate the solvent. Purify using SCX ion-exchange chromatography. The solid obtained was submitted to HPLC purification (the purification was carried out using basic conditions with 60% of Am. Bicarbonate at pH 8 and 40 % of acetonitrile) to afford the title compound (5.5 mg, 43%).


$^1\text{H-NMR}$ (CD_3OD , 300 MHz): 8.62 (s, 1H), 8.11-8.05 (m, 2H), 7.68-7.55 (m, 3H), 4.02 (s, 2H), 2.78-2.73 (m, 2H), 1.72-1.59 (m, 1H), 1.51 (q, 2H, $J = 7.0$ Hz), 0.94 (d, 6H, $J = 6.6$ Hz). Electrospray MS $M+1$ ion = 311.

Example 36

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide

Step 1

6-Amino-5-bromo-nicotinonitrile




To a solution of 6-Amino-nicotinonitrile (102 mg, 0.86 mmol) in AcOH (2 mL) add a solution of Br_2 1.0M in AcOH (0.86 mmol). Stir the mixture at room temperature for 2 hours. Eliminate the solvent. Purify by flash chromatography on silica gel (eluent: hexane/EtOAc 3/1) to afford the title compound (110 mg, 65%).

$^1\text{H-NMR}$ (DMSO, 300 MHz): 8.26 (d, 1H, $J = 2.0$ Hz), 8.10 (d, 1H, $J = 2.0$ Hz), 7.24 (bs, 2H). MS (APCI Neg): 196, 198

Step 2


5-Bromo-6-chloro-nicotinonitrile



To a solution of anhydrous CuCl_2 (77 mg, 0.58 mmol) in CH_3CN (3 mL) add tert-BuONO (0.72 mmol). Heat the mixture at 65°C and then add a suspension of the intermediate above (96 mg, 0.48 mmol) in CH_3CN (2 mL). Stir the mixture for 3 hours. Cool at room temperature. Pour into HCl 3M and extract with EtOAc . Dry the organic layer over Na_2SO_4 . Eliminate the solvent. Purify by flash chromatography on silica gel (eluent: hexane/ EtOAc 2/1) to afford the title compound (55 mg, 55%).
 $^1\text{H-NMR}$ (DMSO , 300 MHz): 8.93 (s, 1H), 8.90 (s, 1H).

Step 3


2-Bromo-4-[(3-methyl-butylamino)-methyl]-phenol



Combine 3-Bromo-4-hydroxy-benzaldehyde (1.5 g, 7.5 mmol) with 3-methylbutylamine (867 μL , 7.5 mmol), MeOH (25 mL) and molecular sieves 3A (5.8 g) and stir at room temperature overnight. Add NaBH_4 (1.4 g, 37.3 mmol) and stir at room temperature for 2 hours. Filtrate over celite with MeOH and eliminate the solvent. Purify by SCX ion-exchange chromatography to afford the title compound (1.5 g, 75%).
 $^1\text{H-NMR}$ (CD_3OD , 300 MHz): 7.52 (d, 1H, $J = 2.0$ Hz), 7.17 (dd, 1H, $J = 2.0$ and 8.5 Hz), 6.87 (d, 1H, $J = 8.5$ Hz), 3.80 (s, 2H), 2.79-2.74 (m, 2H), 1.76-1.63 (m, 1H), 1.54 (q, 2H, $J = 6.8$ Hz), 0.99 (d, 6H, $J = 6.56$ Hz).
 $^{13}\text{C-NMR}$ (CD_3OD , 300 MHz): 158.6, 134.9, 130.7, 128.6, 118.8, 112.7, 53.3, 47.9, 38.6, 27.8, 23.3.

Step 4


(3-Bromo-4-hydroxy-benzyl)-(3-methyl-butyl)-carbamic acid tert-butyl ester



Dissolve intermediate above (800 mg, 2.94 mmol) in dry THF (5 mL) under N_2 atmosphere. Add a solution of di-tert-butyl dicarbonate (642 mg, 2.94 mmol) in dry THF (4 mL). Stir the mixture at room temperature overnight. Eliminate the solvent. Purify by ISCO chromatography (eluent: EtOAc/hexane 1/5) to obtain the title compound (850 mg, 78%). 1H -NMR ($CDCl_3$, 300 MHz): 7.34 (s, 1H), 7.07 (bs, 1H), 6.96 (d, 1H; J = 8.1 Hz), 4.13 (bs, 2H), 3.12 (bs, 2H), 1.48-1.24 (m, 12H), 0.88 (d, 6H, J = 6.6 Hz). Electrospray MS M-1 ion= 370, 372.

Step 5


3-Bromo-4-(3-bromo-5-cyano-pyridin-2-yloxy)-benzyl]-(3-methyl-butyl)-carbamic acid tert-butyl ester



Heat a mixture of the phenol obtained in step 4 (677 mg, 1.82 mmol), 5-Bromo-6-chloro-nicotinonitrile (395 mg, 1.82 mmol), K_2CO_3 (277 mg, 2.0 mmol) and DMF (22 mL) at $100^\circ C$ under N_2 atmosphere overnight. Cool at room temperature. Pour into ice-water and extract with EtOAc. Dry the organic layer over Na_2SO_4 . Eliminate the solvent. Purify by flash chromatography on silica gel (eluent: hexane/EtOAc 8/1) to afford the title compound (860 mg, 85%). 1H -NMR ($CDCl_3$, 300 MHz): 8.30 (d, 1H, J = 2.0 Hz), 8.19 (d, 1H, J = 2.0 Hz), 7.53 (s, 1H), 7.29 (m, 1H), 7.18 (d, 1H, J = 8.3 Hz), 4.43 (m, 2H), 3.19 (m, 2H), 1.58-1.42 (m, 12H), 0.90 (d, 6H, J = 6.5 Hz).

Step 6

[4-(5-Cyano-3-vinyl-pyridin-2-yloxy)-3-vinyl-benzyl]-(3-methyl-butyl)-carbamic acid tert-butyl ester




Heat a mixture of the intermediate above (154 mg, 0.28 mmol), $PdCl_2(dppf)$ (23 mg), tributyl-vinyl-stannane (1.67 mmol) in dry DMF under N_2 atmosphere at $125^\circ C$ for 5

hours. Cool at room temperature. Filtrate over celite with EtOAc. Wash the organic layer with saturated NaCl and then saturated potassium fluoride (KF). Dry the organic layer over Na₂SO₄. Eliminate the solvent. Purify by flash chromatography on silica gel (eluent: hexane/EtOAc 40/1, 20/1 and 10/1) to afford the title compound (70 mg, 58%). ¹H-NMR (CDCl₃, 300 MHz): 8.25 (d, 1H, J= 2.0 Hz), 8.02 (d, 1H, J= 2.4 Hz), 7.49 (bs, 1H), 7.19 (bs, 1H), 7.07-6.97 (m, 2H), 6.64 (dd, 1H, J= 10.9 and 17.8 Hz), 6.01 (d, 1H, J= 17.4 Hz), 5.72 (dd, 1H, J= 0.8 and 17.8 Hz), 5.61 (d, 1H, J= 11.3 Hz), 5.23 (d, 1H, J= 11.7 Hz), 4.44 (bs, 2H), 3.1 (bs, 2H), 1.48 (bs, 12H), 0.89 (d, 6H, J= 6.5 Hz). Electrospray MS M+1-^tBu ion= 392.

Step 7


(2-Cyano-5-oxa-4-aza-dibenzo[a,d]cyclohepten-8-ylmethyl)-(3-methyl-butyl)-carbamic acid tert-butyl ester



Dissolve intermediate above (142 mg, 0.32 mmol) in 1,2-DCE (previously dried over molecular sieves 3A, 33 mL), add Grubb's 2nd generation catalyst (54 mg, 0.06 mol). Heat the mixture under N₂ atmosphere at 82°C during 6 hours. Cool at room temperature. Eliminate the solvent. Purify the by ISCO chromatography (eluent: hexane/EtOAc 8%-20%) to afford the title compound (60 mg, 46%). ¹H-NMR (CDCl₃, 300 MHz): 8.53 (d, 1H, J= 2.0 Hz), 7.78 (d, 1H, J= 2.0 Hz), 7.31-7.17 (m, 2H), 7.06 (bs, 1H), 6.82 (d, 1H, J= 11.5 Hz), 6.53 (d, 1H, J= 11.3 Hz), 4.37 (s, 2H), 3.13 (bs, 2H), 1.49-1.37 (bs, 12H), 0.88 (d, 6H, J= 6.5 Hz). Electrospray MS M+1 ion= 420.

Step 8


(2-Carbamoyl-5-oxa-4-aza-dibenzo[a,d]cyclohepten-8-ylmethyl)-(3-methyl-butyl)-carbamic acid tert-butyl ester



To a solution of the intermediate above (50 mg, 0.12 mmol) in dry DMSO (1.5 mL) add K_2CO_3 (8.2 mg, 0.06 mmol), cool the mixture at 0°C and add H_2O_2 33% (0.48 mmol) dropwise. Stir the mixture at room temperature overnight. Pour into ice-water and extract with EtOAc, dry the organic layer over Na_2SO_4 . Eliminate the solvent at reduced pressure. Purify by ISCO chromatography (eluent: EtOAc) to obtain the title compound (39 mg, 78%). 1H -NMR (MeOD, 300 MHz): 8.53 (d, 1H, J = 2.0 Hz), 7.78 (d, 1H, J = 2.0 Hz), 7.31-7.17 (m, 2H), 7.06 (bs, 1H), 6.82 (d, 1H, J = 11.5 Hz), 6.53 (d, 1H, J = 11.3 Hz), 4.37 (s, 2H), 3.13 (bs, 2H), 1.49-1.37 (bs, 12H), 0.88 (d, 6H, J = 6.5 Hz).

Step 9


(2-Carbamoyl-10,11-dihydro-5-oxa-4-aza-dibenzo[a,d]cyclohepten-8-ylmethyl)-(3-methyl-butyl)-carbamic acid tert-butyl ester



Stir a mixture of the intermediate from Step 8 Example 35 (38 mg, 0.09 mmol) and Pd/C 10% (4 mg) under H_2 atmosphere (1 atm) during 48h. Filter over celite with MeOH. Eliminate the solvent to afford the title compound (35 mg, 92%). 1H -NMR (MeOD, 300 MHz): 8.62 (m, 1H), 8.17 (m, 1H), 7.26-7.13 (m, 3H), 4.42 (s, 2H), 3.24 (m, 2H), 3.18 (bs, 4H), 1.51-1.38 (m, 12H), 0.89 (d, 6H, J = 6.5 Hz).

Step 10

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide




Dissolve the intermediate from step 10 above (30 mg, 0.07 mmol) in CH_2Cl_2 (1.5 mL). Add trifluoroacetic acid (1.82 mmol) and stir the mixture at room temperature overnight. Eliminate the solvent. Submit the crude to a SCX ion-exchange chromatography to afford a yellow oil. Purify by HPLC (basic conditions with Ammonium Bicarbonate (10 mM) at

pH 8. Isocratic mode: 27 % CH₃CN) to afford the title compound (20 mg, 67%). ¹H-NMR (MeOD, 300 MHz): 8.61 (d, 1H, J= 1.9 Hz), 8.17 (d, 1H, J= 2.0 Hz), 7.28-7.26 (m, 3H), 3.80 (s, 2H), 3.19 (s, 4H), 2.70-2.65 (m, 2H), 1.70-1.57 (m, 1H), 1.50-1.43 (q, 2H, J= 6.8 Hz), 0.92 (d, 6H, J= 6.7 Hz). Electrospray MS M+1 ion= 340.

Example 37


8-[(3-Methyl-butylamino)-methyl]-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide



Dissolve the intermediate obtained in step 8 of example 36 (9 mg) in CH₂Cl₂ (1 mL). Add trifluoroacetic acid (50 μl) and stir the mixture at room temperature overnight. Eliminate the solvent. Purify by flash chromatography (eluent: eluent: CHCl₃/B 10%; B: EtOH/NH₄OH 10%) to afford the title compound (3.5 mg, 51%). ¹H-NMR (MeOD, 300 MHz): 8.70 (d, 1H, J= 2.2 Hz), 8.20 (d, 1H, J= 2.2 Hz), 7.42-7.27 (m, 3H), 6.91 (d, 1H, J= 11.3 Hz), 6.75 (d, 1H, J= 11.4 Hz), 3.78 (s, 2H), 2.66-2.61 (m, 2H), 1.69-1.56 (m, 1H), 1.45 (q, 2H, J= 6.8 Hz), 0.91 (d, 6H, J= 6.7 Hz). Electrospray MS M+1 ion= 338.

CLAIMS

1. A compound of formula (I)



j is 1 or 2;

y is 0, 1, or 2; and z is 0, 1, or 2;

p is 0, 1, or 2;

wherein E is O or NH; and wherein each of

X₁, X₂, X₃, X₄, X₅, or X₆, is C, CH, or N; provided that each of rings A or C has no more than 2 nitrogen atoms; and provided that Ring B has 0 or 1 double bond;

R¹ and R² are independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, C₁-C₁₀ alkylaryl, C(O)C₁-C₈ alkyl, SO₂C₁-C₈alkyl, SO₂C₁-C₈alkylNR⁸R⁸, (CH₂)_nC(O)NR⁸R⁸, SO₂C₁-C₁₀ alkylaryl, SO₂C₁-C₈ alkylheterocyclic, C₄-C₁₀ alkylcycloalkane, (CH₂)_nC(O)OR⁸, and (CH₂)_nC(O)R⁸; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C₁-C₈ alkyl, C₂-C₈ alkenyl, phenyl, C₃-C₈ cycloalkyl, C₁-C₈ alkylaryl, and C(O)C₁-C₈ alkyl; and wherein R¹ and R² may optionally combine with each other to form a 4, 5, 6, or 7-membered nitrogen-containing heterocycle which nitrogen-containing heterocycle may further have substituents selected from the group consisting of oxo, amino, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, C₁-C₈ alkylaryl, C(O)C₁-C₈ alkyl, CO(O)C₁-C₈ alkyl, halo, C₁-C₈ haloalkyl;

R³ and R^{3'} are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, phenyl, aryl, C₁-C₈ alkylcycloalkyl, and C₁-C₈ alkylaryl;

R^a and R^b are each independently selected from hydrogen, and C₁-C₃ alkyl or combine with their respective carbon atoms to form the vinyl diradical -CH=CH-;

R⁴ and R⁵ are each independently selected from hydrogen, C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, C₁-C₈ alkoxy, halo, C₁-C₈ haloalkyl, phenyl, aryl, C₁-C₈ alkylaryl, (CH₂)_mNSO₂C₁-C₈ alkyl, (CH₂)_mNSO₂phenyl, (CH₂)_mNSO₂aryl, -C(O)C₁-C₈ alkyl, and -

C(O)OC_{1-C8} alkyl; wherein each R⁴ and R⁵ is attached to its respective ring only at carbon atoms; wherein m is 1 or 2; and n is 1, 2, or 3;

R⁶ and R⁷ are each independently selected from hydrogen, C_{1-C8} alkyl, C_{2-C8} alkenyl, C_{2-C8} alkynyl, C(O)C_{1-C8} alkyl, SO₂C_{1-C8} alkyl, SO₂C_{1-C8} alkylaryl, SO₂C_{1-C8} alkylheterocyclic, aryl, C_{1-C8} alkylaryl, C_{3-C7} cycloalkane, C_{1-C8} alkylcycloalkane, (CH₂)_nC(O)OR⁸, (CH₂)_nC(O)R⁸, (CH₂)_mC(O)NR⁸R⁸, and (CH₂)_mNSO₂R⁸; wherein each of the alkyl, alkenyl, and aryl groups are optionally substituted with one to five groups independently selected from C_{1-C8} alkyl, C_{2-C8} alkenyl, phenyl, and C_{1-C8} alkylaryl; and wherein R⁶ and R⁷ may independently combine with each other, and with the nitrogen atom to which they are attached to form a 4, 5, 6, or 7-membered nitrogen containing heterocycle which nitrogen containing heterocycle may optionally have substituents selected from the group consisting of oxo, amino, C_{1-C8} alkyl, C_{2-C8} alkenyl, C_{2-C8} alkynyl, phenyl, and C_{1-C8} alkylaryl;

R⁸ is independently selected from hydrogen, C_{1-C8} alkyl, C_{2-C8} alkenyl, phenyl, benzyl, and C_{5-C8} alkylaryl; or a pharmaceutically acceptable salt, solvate, prodrug, tautomers, enantiomer, racemate, diastereomer, or mixture of diastereomers thereof.

2. The compound according to claim 1 wherein the A-ring is selected from the group consisting of phenyl, pyridine, pyrimidine and pyrazine.

3. A compound according to Claim 1 wherein the C-ring is selected from the group consisting of phenyl and pyridine.

4. A compound according to Claim 1 wherein the A-ring is phenyl and the C ring is pyridine.

5. A compound according to Claim 1 wherein both A and C rings are phenyl.

6. A compound according to claim 1 wherein p is 2 and both R^a and R^b are hydrogen.

7. A compound according to Claim 1 wherein -(CR^aR^b)_p- equals -CH=CH-.

8. A compound according to Claim 1 wherein E is an oxygen atom.

5

10

[illegible]

15

20

14. A compound selected from the group consisting of:
8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid
amide,
8-(Isobutylamino-methyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(4-Methyl-pentylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(2-Thiophen-2-yl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Pentylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Hexylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(Cyclohexylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Cyclooctylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-Cycloheptylaminomethyl-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(Cycloheptylmethyl-amino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide trifluoroacetate salt,

8-[(2-(Tetrahydro-pyran-4-yl)-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3,3-Dimethyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(2-Cyclopentyl-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Morpholin-4-yl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Ethoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(2-Diethylamino-ethylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Methoxy-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide, and

8-[(3-Phenyl-propylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(3-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(3-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[2-(4-Chloro-phenyl)-pyrrolidin-1-ylmethyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Phenyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Phenyl-piperidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Phenyl-azepan-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-(2-Benzyl-pyrrolidin-1-ylmethyl)-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-dibenzofuran-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-dibenzo[b,f]oxepine-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-10,11-dihydro-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide,

8-[(3-Methyl-butylamino)-methyl]-5-oxa-4-aza-dibenzo[a,d]cycloheptene-2-carboxylic acid amide,

or a pharmaceutically acceptable salt, solvate, enantiomer, diastereomer and diastereomeric mixture thereof.

15. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of Claims 1 to 14 in association with a carrier, diluent and/or excipient.

16. A method for blocking a mu, kappa, delta or receptor combination (heterodimer) thereof in mammals comprising administering to a mammal requiring blocking of a mu, kappa, delta or receptor combination (heterodimer) thereof, a receptor blocking dose of a compound according to any one of Claims 1 to 14, or a pharmaceutically acceptable salt, enantiomer, racemate, mixture of diastereomers, or solvate thereof.

17. A method of treating or preventing obesity and Related Diseases comprising administering a therapeutically effective amount of a compound of formula I.

18. A method according to Claim 16 wherein the Related Diseases is selected from the group consisting of diabetes, diabetic complications, diabetic retinopathy,

atherosclerosis, hyperlipidemia, hypertriglycemia, hyperglycemia, and hyperlipoproteinemia.


5 19. A method of treating and/or preventing diseases related to obesity including irritable bowel syndrome, nausea, vomiting, depression, smoking and alcohol addiction, sexual dysfunction, substance abuse, drug overdose, addictive behavior disorders, compulsive behaviors, and stroke comprising administering a therapeutically effective amount of a compound of formula I.

10 20. A method of suppressing appetite in a patient in need thereof, comprising administering a therapeutically effective amount of a compound of formula I.

15 21. Use of a compound of formula I in the manufacture of a medicament for the treatment and/or amelioration of the symptoms associated with obesity and Related Diseases comprising administering a therapeutically effective amount of a compound of formula I to a patient in need thereof.

ABSTRACT

A compound of the formula (I)



(I)

wherein the variables X₁ to X₆, R^a, R^b, R¹ to R⁷ including R³, E, p, j, y, z, A, B and C are as described or a pharmaceutically acceptable salt, solvate, enantiomer, racemate, diastereomer or mixtures thereof, useful for the treatment, prevention or amelioration of obesity and Related Diseases is disclosed.

